

LOT NUMBER AND EXPIRATION DATE SHALL APPEAR ABOVE THE WEBSITE

NON-VARNISH AREA

BARCODE TYPE C128
TO READ U5530R

U5530R

Open Here

Visit www.primatene.com
See Insert and Side Panels
For Special Directions On:
■ Activating Your New Inhaler
■ Dosing with Your New Inhaler
■ Using Spray Indicator

Drug Facts

Active ingredient Purpose
(in each spray)
Epinephrine 0.125 mg.....Bronchodilator

Uses
for temporary relief of mild symptoms of intermittent asthma
■ wheezing
■ tightness of chest
■ shortness of breath

Warnings
For oral inhalation only
Asthma alert: Because asthma may be life threatening, see a doctor if you
■ are not better in 20 minutes
■ get worse
■ need more than 8 inhalations in 24 hours
■ have more than 2 asthma attacks in a week
These may be signs that your asthma is getting worse.
Do not use
■ unless a doctor said you have asthma
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs taken for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or a pharmacist before taking this product.

Ask a doctor before use if you have
■ ever been hospitalized for asthma
■ heart disease
■ high blood pressure
■ diabetes
■ trouble urinating due to an enlarged prostate gland
■ thyroid disease
■ seizures
■ narrow angle glaucoma

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are
■ taking prescription drugs for asthma, obesity, weight control, depression, or psychiatric or emotional conditions
■ taking any drug that contains phenylephrine, pseudoephedrine, ephedrine, or caffeine (such as for allergy, cough-cold, or pain)

When using this product
■ your blood pressure or heart rate may go up. This could increase your risk of heart attack or stroke, which may cause death.
■ your risk of heart attack or stroke increases if you:
■ have a history of high blood pressure or heart disease
■ take this product more frequently or take more than the recommended dose.
■ avoid foods or beverages that contain caffeine
■ avoid dietary supplements containing ingredients reported or claimed to have a stimulant effect
■ avoid spraying in eyes
■ contents under pressure. Do not puncture or incinerate.
■ do not store near open flame or heat above 120°F (49°C). May cause bursting.

Stop use and ask a doctor if
■ your asthma is getting worse (see asthma alert)
■ you have difficulty sleeping
■ you have a rapid heart beat
■ you have tremors, nervousness, or seizure

If pregnant or breast-feeding, ask a health care professional before use
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ read the Consumer Information Insert for detailed directions on how to use this product
■ do not use more than directed

Drug Facts (continued)

■ for adults and children 12 years of age and over
■ children under 12 years of age: do not use; it is not known if the drug works or is safe in children under 12

Before first use (new inhaler)
activate new inhaler by shaking then spraying into air 4 separate times
each time you dose
■ remove red cap (if attached)
■ shake then spray into the air 1 time
■ exhale completely, place inhaler in mouth
■ inhale deeply while pressing down on top of inhaler, then continue the deep breath
■ hold breath as long as possible, exhale
■ wait 1 minute. If symptoms not relieved, take a second inhalation by repeating steps above.

after use
■ wait at least 4 hours between doses
■ do not use more than 8 inhalations in 24 hours
■ wash inhaler after each day of use. Run water through the mouthpiece for 30 seconds.

Other information
■ store at room temperature, between 15-25°C (59-77°F)
■ contains no sulfites
■ keep this label and enclosed materials. They contain important additional information.

Inactive ingredients
dehydrated alcohol (1%), hydrofluoroalkane (HFA 134a), polysorbate 80, thymol

Questions or comments?
call 1-8 Primatene or 1-877-462-8363
Mon.- Fri. 7 am - 5 pm PST

Primatene[®]
MIST

Epinephrine Inhalation Aerosol
0.125 mg per spray
Bronchodilator

For Oral
Inhalation Only

For **TEMPORARY** relief
of **MILD** symptoms
of **INTERMITTENT** asthma



Suspension:
Shake then Spray
into the air before
each inhalation.

NEW FORMULATION:
See Important Usage
Information on Insert
and on Side Panels

160 Metered Sprays
Net Weight: 11.7g

UPC-A BARCODE
TO READ: 3 17270 55300 7

Manufactured by:
Armstrong Pharmaceuticals, Inc.
Canton, MA 02021
© 2018 Made in USA

NON-VARNISH AREA

BARCODE TYPE C128
TO READ U5530R

U5530R

DESCRIPTION: Primatene MIST
LABEL NO: U5530R
SIZE: 1.9375 X 1.706 X 5.125
COLORS: PMS 4625, PMS 485, PMS 430, PMS 7406 C, CMYK
BARCODES: CODE 128 TO READ: U5530R
UPC-A TO READ: 317270553007